

# Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector



Circumvent<sup>®</sup> G2  
PCV

**500 mL**

**Option 1:**  
250 Doses-2 mL

**Option 2:**  
500 Doses-1 mL



For use in healthy swine as an aid in the prevention of viremia, aid in the reduction of virus shedding and aid in the reduction of lymphoid infection caused by PCV 2. Efficacy for at least 20 weeks following completion of vaccination has been demonstrated.

**ADMINISTRATION and DOSAGE OPTIONS:** Allow the vaccine to reach an ambient temperature of 16°- 27° C (60°- 80° F) and shake well prior to use. Use aseptic technique. **OPTION 1:** Vaccinate pigs intramuscularly at 3 weeks of age or older with a single 2 mL dose. **OPTION 2:** Vaccinate pigs intramuscularly as early as 3 days of age with a 1 mL dose followed by a second 1 mL dose 3 weeks later.

**CAUTIONS:** Store refrigerated at 2°-7°C (35°-45°F). Do not freeze. Use entire contents when first opened; do not save partial contents. Burn the container and all unused product. Use only in healthy swine. Do not vaccinate within 21 days before slaughter. The vaccine may induce significant localized adverse reactions, particularly in pigs following booster vaccination. Localized reactions resolve 1-35 days following vaccination. If allergic reaction occurs, treat with epinephrine. Caution should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. Contact a physician immediately.

**PRESERVATIVES:** Gentamicin and thimerosal.  
**FOR ANIMAL USE ONLY**

Intervet Inc., Merck Animal Health, division of Intervet Inc.  
Omaha, Nebraska 68103 USA  
U.S. Vet Lic. No. 165A • 1 800 521-5767



Code 119100  
135844-03

Serial No.  
Exp. Date